Asthma Action Plan



General Information:				
■ Name Name Emergency contact ■ Physician/healthcare provider			Phone numbersPhone numbers	
Severity Classification O Intermittent O Moderate Persistent O Mild Persistent O Severe Persistent	Triggers Colds O Smoke O Weather Exercise O Dust O Air Pollution Animals O Food Other		1. Premedication (how much and when) 2. Exercise modifications	
Green Zone: Doing Well	Peak Flow Meter Pe	rsonal Best =		
Symptoms	Control Medications:			
■ Breathing is good ■ No cough or wheeze ■ Can work and play ■ Sleeps well at night Peak Flow Meter More than 80% of personal best or	1			When to Take It
Yellow Zone: Getting Worse Symptoms Some problems breathing Cough, wheeze, or chest tight Problems working or playing	Contact physician if using quick rel Continue control medicines and add: Medicine How Much to Ta		ake	When to Take It
Wake at night Peak Flow Meter Between 50% and 80% of personal best orto	IF your symptoms (and peak flow, if used) return to Green Zone after one hour of the quick-relief treatment, THEN		IF your sym	nptoms (and peak flow, if used) urn to Green Zone after one quick-relief treatment, THEN
	 Take quick-relief medication every 4 hours for 1 to 2 days. Change your long-term control medicine by 		 Take quick-relief treatment again. Change your long-term control medicine by Call your physician/Healthcare provider 	
	O Contact your physician fo		medication	hour(s) of modifying your on routine.
Red Zone: Medical Alert	Ambulance/Emergency Phone Number:			
Symptoms ■ Lots of problems breathing ■ Cannot work or play ■ Getting worse instead of better ■ Medicine is not helping	Medicine	rol medicines and add: How Much to Ta		When to Take It
Peak Flow Meter Less than 50% of personal best or to	 Go to the hospital or call for an ambulance if: Still in the red zone after 15 minutes. You have not been able to reach your physician/healthcare provider for help. 		following o	danger signs are present: walking/talking due to shortness

O Lips or fingernails are blue.